

Horizons in Robinson-Trautman space-times

W. Natorf, J. Tafel

Institute of Theoretical Physics, Warsaw University,
ul. Hoża 69, 00-681 Warsaw, Poland
e-mail: nat@fuw.edu.pl, tafel@fuw.edu.pl

Abstract

The past quasi-local horizons in vacuum Robinson-Trautman space-times are described. The case of a null (non-expanding) horizon is discussed. It is shown that the only Robinson-Trautman space-time admitting such a horizon with sections diffeomorphic to S_2 is the Schwarzschild space-time. Weakening this condition leads to the horizons of the C-metric. Properties of the hypersurface $r = 2m$ are examined.

1 Introduction

A vacuum Robinson-Trautman (RT) metric can be determined by a function of three variables, satisfying a fourth order partial differential equation [1]. Very few of its solutions are known in an explicit form [2]. The only ones representing asymptotically flat space-times correspond to the Minkowski and the Schwarzschild metrics. Nontrivial asymptotically flat RT space-times were shown to exist and tend to the Schwarzschild solution in the limit of infinite retarded time [3].

In this paper we describe quasi-local horizons [4] in RT space-times. We derive equations obtained previously by Tod [5] and Chow and Lun [6]. Using the equation of Chow and Lun we prove that the past horizon is null if and only if its sections admit a shear-free inward pointing null normal vector field.

It is a well known property of a nonexpanding horizon that its null generator is the repeated principal null direction of the Weyl tensor [4]. Since the repeated principal null direction responsible for algebraic properties of the Weyl tensor of RT space-time is transverse to the horizon, the Weyl tensor is of Petrov type D on the horizon. We prove that this property holds also in the neighbourhood of the horizon and no nonexpanding horizons of the form $R \times S_2$ exist unless the RT space-time is the Schwarzschild space-time. If no restriction on topology of sections is made, one obtains null horizons of the C-metric. Their sections contain conical singularities [7].

In section 5 we consider the metric induced on the hypersurface $r = 2m$ for finite u , and examine its signature (Proposition 5.2). We also find the relation between this hypersurface and trapped surfaces (Proposition 5.1).

The standard form of the Robinson-Trautman metric reads [1, 2]

$$g = 2du(Hdu + dr) - 2\frac{r^2}{P^2}d\xi d\bar{\xi} . \quad (1)$$

Vacuum Einstein's equations give

$$H = P^2(\ln P)_{,\xi\bar{\xi}} - r(\ln P)_{,u} - \frac{m}{r} \quad (2)$$

and the Robinson-Trautman equation

$$P^2(P^2(\ln P)_{,\xi\bar{\xi}})_{,\xi\bar{\xi}} + 3m(\ln P)_{,u} - m_{,u} = 0 . \quad (3)$$

Here m is a function of u which, when non-vanishing, can be transformed to $m = \pm 1$. In this case equation (3) takes the form

$$K_{,\xi\bar{\xi}} - 3m(P^{-2})_{,u} = 0 \quad (4)$$

where

$$K = 2P^2(\ln P)_{,\xi\bar{\xi}} \quad (5)$$

is the Gaussian curvature of the surface given by $u = \text{const}$, $r = 1$. Point transformations preserving metric (1) and m have the following form [2]:

$$\begin{aligned} u &\rightarrow u + \text{const} , \\ \xi &\rightarrow h(\xi) , \\ P &\rightarrow |h'|^{-1}P , \\ r &\rightarrow r . \end{aligned} \quad (6)$$

The induced metric of a surface of constant u and r is given by

$$g' = 2\frac{r^2}{P^2}d\xi d\bar{\xi} . \quad (7)$$

If the function

$$\hat{P} = P/(1 + \xi\bar{\xi}/2) \quad (8)$$

is regular (smooth and positive) for all $\xi \in C$ and $\xi \rightarrow \infty$, then the surface $u = \text{const}$, $r = \text{const}$ is diffeomorphic to S_2 and ξ can be interpreted as the complex stereographic coordinate on S_2 . The function $P = (1 + \xi\bar{\xi}/2)$ giving the standard metric of S_2 will be denoted by P_S . Two dimensional surfaces admitting regular $\hat{P}$ will be referred to as regular surfaces.

2 Marginally trapped surfaces

Let $\mathcal{S}$ be a 2-dimensional space-like surface given by

$$u = \text{const}, \quad r = R(\xi, \bar{\xi}) . \quad (9)$$

We introduce four null forms,

$$\begin{aligned} \theta^0 &= du , \\ \theta^1 &= \left(H + \frac{P^2}{r^2} |R_{,\xi}|^2 \right) du + dr - (R_{,\xi} d\xi + \text{c.c.}) , \\ \theta^2 &= \frac{r}{P} d\xi - \frac{P}{r} R_{,\bar{\xi}} du , \quad \theta^3 = \bar{\theta}^2 , \end{aligned} \quad (10)$$

and the dual null tetrad,

$$\begin{aligned} e_0 &= \partial_u - \left(H - \frac{P^2}{r^2} |R_{,\xi}|^2 \right) \partial_r + \frac{P^2}{r^2} (R_{,\xi} \partial_\xi + \text{c.c.}) , \\ e_1 &= \partial_r , \\ e_2 &= \frac{P}{r} (\partial_\xi + R_{,\xi} \partial_r) , \quad e_3 = \bar{e}_2 . \end{aligned} \quad (11)$$

In Appendix A we list the non-vanishing Newman-Penrose spin coefficients and Weyl scalars related to the tetrad (11), needed in further calculations. The ingoing and outgoing null vectors normal to $\mathcal{S}$ are, respectively, $l = e_0$, $k = e_1$, while $m = e_2$ and $\bar{m} = e_3$ are tangent to $\mathcal{S}$.

Using the formulae

$$\begin{aligned} -m^a \bar{m}^b l_{(a;b)} &= \frac{H}{r} + (\ln P)_{,u} - \frac{P^2}{r^2} R_{,\xi\bar{\xi}} + \frac{P^2}{r^3} |R_{,\xi}|^2 , \\ -m^a \bar{m}^b k_{(a;b)} &= -r^{-1} \end{aligned} \quad (12)$$

from Appendix A and substituting for H from (2) we find the expansion scalars of the vectors k and l on $\mathcal{S}$ [5]

$$\theta_{(l)} = \frac{1}{R} \left(\frac{K}{2} - P^2 (\ln R)_{,\xi\bar{\xi}} - \frac{m}{R} \right) , \quad (13)$$

$$\theta_{(k)} = -R^{-1} . \quad (14)$$

Therefore, the condition that the expansion of l vanishes takes the form of the following equation for R :

$$-P^2 (\ln R)_{,\xi\bar{\xi}} + \frac{K}{2} - \frac{m}{R} = 0 . \quad (15)$$

The existence and uniqueness of positive, smooth solutions of (15) for regular spherical surfaces $u = \text{const}$, $r = \text{const}$ has been proved by Tod [5]. If $m = 1$, the area of these surfaces is independent of u due to (3) and Stokes' theorem, and can be chosen to be equal to 4π . Note that u enters (15) as a parameter via the function P .

3 Horizons

In this section we consider hypersurfaces foliated by marginally trapped surfaces. Let $\mathcal{H}$ be defined by

$$r = R(u, \xi, \bar{\xi}) , \quad (16)$$

where R satisfies (15). It follows that

$$\nabla_t(m^a \bar{m}^b l_{(a|b)}) = 0 \quad (17)$$

on $\mathcal{H}$, where t is tangent to $\mathcal{H}$ and orthogonal to its sections. Using maximum principle and (13) Tod proved that a marginally trapped surface described by (9) and (15) is outermost [5]. Results of Andersson et al. [8] assure that there exists a horizon such that $\mathcal{S}$ is its section.

Using (11) we obtain

$$t = l + \chi k , \quad \chi = R_{,u} + H + \frac{P^2}{R^2} R_{,\xi} R_{,\bar{\xi}} . \quad (18)$$

Following [6] we rewrite equation (17) in terms of the spin coefficients:

$$\frac{P^2}{R} \left(\frac{\chi}{R} \right)_{,\xi \bar{\xi}} + \chi \Psi_2 - \lambda \bar{\lambda} = 0 , \quad (19)$$

where λ is the shear of l . Equation (19) corresponds to equation (15) in [9], where the general case of constraint equations on a marginally trapped tube was considered.

Assume now that the surfaces $u = \text{const}$, $r = \text{const}$ have spherical topology. As pointed out in [6], it follows from the maximum principle for (19) that $\mathcal{H}$ is a non-timelike surface. If $\mathcal{H}$ is null ($\chi = 0$), then $\lambda = 0$. Conversely, if λ vanishes, multiplying both sides of (19) by χ/R^2 and using Stokes' theorem for $\mathcal{S}$ we get

$$- \int_{\mathcal{S}} \left| \nabla \frac{\chi}{R} \right|^2 d\sigma = \int_{\mathcal{S}} \frac{\chi^2}{R^2} |\Psi_2| d\sigma , \quad (20)$$

where $d\sigma = i(P/R)^2 d\xi \wedge d\bar{\xi}$ is the natural measure on $\mathcal{S}$ and $\Psi_2 = -|\Psi_2|$ follows from (A.9) and the assumption $m > 0$. This shows that χ vanishes. Thus,

$$\mathcal{H} \text{ is null} \Leftrightarrow \lambda = 0 . \quad (21)$$

4 Null case

We would like to discuss conditions that R and P should satisfy for $\mathcal{H}$ to be a null (nonexpanding) horizon. In this case $\lambda = \chi = \nu = 0$ (see (21) and

(A.11)) and the vector tangent to $\mathcal{H}$ and normal to sections is just l (see (18)). Consider the following Newman-Penrose equations

$$\begin{aligned} \delta\lambda - \bar{\delta}\mu &= (\rho - \bar{\rho})\nu + (\mu - \bar{\mu})\pi + \mu(\alpha + \bar{\beta}) + \\ &+ \lambda(\bar{\alpha} - 3\beta) - \Psi_3 + \Phi_{21} , \end{aligned} \quad (22)$$

$$\begin{aligned} \Delta\lambda - \bar{\delta}\nu &= (\bar{\gamma} - 3\gamma - \mu - \bar{\mu})\lambda + \\ &+ (3\alpha + \bar{\beta} + \pi - \bar{\tau})\nu - \Psi_4 . \end{aligned} \quad (23)$$

Note that derivatives present in (22) and (23) are $\Delta = \nabla_l$, $\delta = \nabla_m$ and $\bar{\delta} = \nabla_{\bar{m}}$. These directions are tangent to $\mathcal{H}$; this allows us to conclude that conditions $\Psi_4 = \Psi_3 = 0$ must be satisfied on $\mathcal{H}$. Hence g is of Petrov type D on $\mathcal{H}$ ($\Psi_2 \neq 0$ for non-vanishing m , see (A.9) in Appendix A). From equation (A.10) in Appendix A it follows that on $\mathcal{H}$

$$\frac{3Pm}{R^4}R_{,\bar{\xi}} + \frac{P}{2R^2}K_{,\bar{\xi}} = 0 . \quad (24)$$

This equation can be easily integrated to give R in terms of K :

$$R = 6m(K + a)^{-1} , \quad (25)$$

where $a = a(u)$ is real. Substituting (25) into the condition $\lambda = 0$ (see (A.4) in App. A) we get

$$(P^2 K_{,\bar{\xi}})_{,\bar{\xi}} = 0 . \quad (26)$$

Note that due to (25), (26) is a condition on P only. Due to the fact that r -dependence of the metric is known, further results hold for arbitrary $r > 0$. It turns out that (26) restricts substantially a set of solutions of the RT equation (3), regardless of whether the trapped surfaces are spherical or not:

Proposition 4.1 *The only vacuum Robinson-Trautman metrics admitting a nonexpanding horizon are the Schwarzschild metric and the C-metric.*

Proof: It follows from (26) that

$$P^2 K_{,\bar{\xi}} = f(u, \xi) . \quad (27)$$

Substituting (5) and (27) into (3) yields

$$(fP^{-2})_{,\xi} = 3m(P^{-2})_{,u} . \quad (28)$$

Applying twice the operator $P^2\partial_{\bar{\xi}} + 2PP_{,\bar{\xi}}$ to (28) we obtain

$$P^2(K_{,\bar{\xi}})^2 = 6m(P^2(\ln P)_{,u\bar{\xi}})_{,\bar{\xi}} . \quad (29)$$

Conditions (26) and (29) assure that the corresponding metric is of Petrov type D [2]. For $f = 0$ equations (27) and (28) lead to $P_{,u} = 0$ and $K = \text{const}$,

hence one obtains the Schwarzschild metric. According to [2], an inspection of Kinnersley's list of type D vacuum solutions of the Einstein equations [10] shows that for $f \neq 0$ solutions of (27) and (28) necessarily yield the C-metrics. We present an independent proof of this property in Appendix B.

In the case of the Schwarzschild metric the trapped surfaces form the past event horizon $r = 2m$. In the case of the C-metric, by virtue of (25), equation (15) takes the form

$$P^2(\ln(K+a))_{,\xi\bar{\xi}} + \frac{K}{3} - \frac{a}{6} = 0 . \quad (30)$$

Substituting (B.18) to (30) and integrating yields

$$6mK_{,x} = -\frac{1}{3}K^3 - \left(\frac{a^2}{3} + a_1\right)K + aa_1 \quad (31)$$

where a_1 is another function of u . Equation (31) coincides with (B.19) if a and a_1 are constants, $a_1 = b - \frac{a^2}{3}$ and

$$\frac{a^3}{3} - ab + c = 0 . \quad (32)$$

Thus, the C-metric admits locally a nonexpanding horizon. It is known that this horizon does not admit regular spherical sections (see e.g. [7]) because of conical singularity on each section $\mathcal{S}$. $\square$

5 Properties of the surface $r = 2m$

It is interesting to examine the relation between $\mathcal{S}$ and the surface $r = 2m$ which plays the role of past event horizon in the Schwarzschild case. In this section we assume that the surfaces of constant r and u are diffeomorphic to S_2 and the function $\hat{P} = P/P_S$ is regular on these surfaces.

Proposition 5.1 *Let $\mathcal{S}$ be a regular spheroidal marginally trapped surface given by (9) and (15). Then $\mathcal{S}$ crosses the surface $r = 2m$.*

Proof: Let

$$R = \frac{2m}{1-h} \quad (33)$$

for some smooth function $h < 1$. Rewriting (15) in terms of h we get

$$-2P^2[\ln(1-h)]_{,\xi\bar{\xi}} = K - 1 + h . \quad (34)$$

From the Gauss-Bonnet theorem and the regularity of h it follows that

$$\int_{\{r=1, u=\text{const}\}} h = 0 , \quad (35)$$

hence $h^{-1}(\{0\}) \neq \emptyset$ which finishes the proof. $\square$

Another property of $r = 2m$ is the signature of its induced metric for finite u . Obviously, it is null in the Schwarzschild space-time. The converse statement is also true:

Proposition 5.2 *Let g be a vacuum RT metric (1). If the hypersurface $r = 2m$ is null, then g is the Schwarzschild metric.*

Proof: Assume that the surface $r = 2m$ is null, then in (1) one has to set $H|_{r=2m} = 0$ which is equivalent to

$$K - 1 = 4m(\ln P)_{,u} . \quad (36)$$

Combining (36) with the RT equation (3) we get

$$\partial_u (4m(\ln P)_{,\xi\bar{\xi}} - 3mP^{-2}) = 0 . \quad (37)$$

Hence

$$(\ln P)_{,\xi\bar{\xi}} - \frac{3}{4}P^{-2} = f(\xi, \bar{\xi}) \quad (38)$$

and

$$K = \frac{3}{2} + 2P^2 f \quad (39)$$

follows. Differentiating K with respect to u and using (36) and (39) we obtain

$$K_{,u} = 4PP_{,u}f = 2(\ln P)_{,u}2P^2f = \frac{1}{2m}(K - 1)(K - 3/2) . \quad (40)$$

This equation can be integrated to give

$$\frac{K - 1}{K - 3/2} = h(\xi, \bar{\xi})e^{-u/4m} \quad (41)$$

for some u -independent function h . Now from (39) and (41) we have

$$K - 1 = he^{-u/4m}2P^2f . \quad (42)$$

Solving (41) for K and using (39) we have

$$P^2 = \frac{1}{4f} \frac{1}{he^{-u/4m} - 1} . \quad (43)$$

Inserting (43) into the RT equation yields a polynomial of degree 4 in $U = \exp(-u/4m)$ equal to zero:

$$(h_{,\xi\bar{\xi}} - 6fh)U - (hh_{,\xi\bar{\xi}} - 2|h_{,\xi}|^2 - 18fh^2)U^2 - 18f(hU)^3 + 6f(hU)^4 = 0 . \quad (44)$$

Since h is u -independent, it has to vanish identically, and from (42) we get $K = 1$ and $P = P_S$. Hence g is the Schwarzschild metric. $\square$

Acknowledgments

The authors would like to thank P. Chruściel and M. Korzyński for helpful comments.

This work was supported by the Polish Ministry of Science and Higher Education grant 1 PO3B 075 29.

Appendix A: Spin coefficients and Weyl scalars for tetrad (10)

We follow the notation from [2] to derive the spin coefficients and Weyl scalars of (10).

$$\alpha = \frac{P_{,\bar{\xi}}}{2r} - \frac{PR_{,\bar{\xi}}}{r^2} , \quad (\text{A.1})$$

$$\beta = -\frac{P_{,\xi}}{2r} , \quad (\text{A.2})$$

$$\tau = \bar{\pi} = -\frac{P}{r^2}R_{,\xi} , \quad (\text{A.3})$$

$$\lambda = (\partial_{\bar{\xi}} + R_{,\bar{\xi}}\partial_r) \left(\frac{P^2}{r^2}R_{,\bar{\xi}} \right) , \quad (\text{A.4})$$

$$\rho = -\frac{1}{r} , \quad (\text{A.5})$$

$$\gamma = \frac{1}{2r^3}[-2P^2|R_{,\xi}|^2 + r^3H_{,r} + rP(P_{,\bar{\xi}}R_{,\xi} - \text{c.c.})] , \quad (\text{A.6})$$

$$\nu = \frac{P}{r}(\partial_{\bar{\xi}} + R_{,\bar{\xi}}\partial_r) \left(R_{,u} + H + \frac{P^2}{r^2}|R_{,\xi}|^2 \right) , \quad (\text{A.7})$$

$$\mu = -\frac{H}{r} - (\ln P)_{,u} + \frac{P^2}{r^2}R_{,\xi\bar{\xi}} - \frac{P^2}{r^3}|R_{,\xi}|^2 , \quad (\text{A.8})$$

$$\Psi_2 = -\frac{m}{r^3} , \quad (\text{A.9})$$

$$\Psi_3 = \frac{3Pm}{r^4}R_{,\bar{\xi}} + \frac{P}{2r^2}K_{,\bar{\xi}} . \quad (\text{A.10})$$

Note that from the definition of χ in (18) and (A.7) it follows that

$$\nu = \nabla_{\bar{m}}\chi . \quad (\text{A.11})$$

Appendix B: Robinson-Trautman metrics of type D

In section 4 we have shown that a necessary condition for $\mathcal{H}$ to be null is that the Robinson-Trautman metric is of type D. In this appendix we will explicitly solve equations (27) and (28) equivalent to the condition $\lambda = 0$ and the Robinson-Trautman equation. Since $f = 0$ corresponds to the Schwarzschild metric or a flat one we will assume here that $f \neq 0$.

Let

$$\xi' = h(u, \xi) \quad (\text{B.1})$$

where

$$h_{,\xi} = f^{-1} \quad (\text{B.2})$$

(note that transformation (B.1) does not preserve the Robinson-Trautman equation (3)). In terms of coordinates ξ' , $\bar{\xi}'$ and $u' = u$ equation (27) reads

$$P^2 K_{,\bar{\xi}'} = |f|^2 . \quad (\text{B.3})$$

Hence, $K = K(u', x')$, where $x' = \text{Re } \xi'$ and

$$P^2 = \frac{2}{K_{,x'} |h_{,\xi}|^2} . \quad (\text{B.4})$$

Substituting (B.4) into definition (5) of K yields

$$(\ln K_{x'})_{,x'x'} = -2KK_{,x'} . \quad (\text{B.5})$$

Integrating twice equation (B.5) we get

$$K_{,x'} = -\frac{1}{3}K^3 + bK + c , \quad (\text{B.6})$$

where b and c are functions of u .

It follows from (B.4) that

$$(\ln P)_{,u} = -\frac{1}{2}(\ln K_{,x'})_{,u'} - \frac{1}{2}(\ln K_{,x'})_{,x'} \text{Re } h_{,u} - (\ln |h_{,\xi}|)_{,u} \quad (\text{B.7})$$

and the Robinson-Trautman equation takes the form

$$(\ln K_{,x'})_{,u'} + (\ln K_{,x'})_{,x'} \left(\text{Re } h_{,u} - \frac{1}{6m} \right) + 2(\ln |h_{,\xi}|)_{,u} = 0 . \quad (\text{B.8})$$

By virtue of (B.6) acting on (B.8) with the operator $\partial_{\xi'} \partial_{\bar{\xi}'} = |h_{,\xi}|^{-2} \partial_{\xi} \partial_{\bar{\xi}}$ yields

$$(\ln (KK_{,x'}))_{,u'} + (\ln (KK_{,x'}))_{,x'} \left(\text{Re } h_{,u} - \frac{1}{6m} \right) + 2(\ln |h_{,\xi}|)_{,u} = 0 \quad (\text{B.9})$$

Subtracting (B.8) from (B.9) implies

$$\frac{K_{,u'}}{K_{,x'}} + \text{Re } h_{,u} - \frac{1}{6m} = 0 . \quad (\text{B.10})$$

Applying $\partial_{\xi'}\partial_{\bar{\xi}'}$ to (B.10) shows that $K_{,u'}/K_{,x'}$ is linear in x' and

$$h_{,u'} = c_1(u)h + c_2(u) \quad (\text{B.11})$$

where c_1 and c_2 are, respectively, a real and a complex function of u . Hence

$$h = \frac{2}{c_3(u)}h'(\xi) + c_4(u) \quad (\text{B.12})$$

where c_3 is a real function of u and h' is a function of ξ . Note that transformation $\xi \rightarrow h'(\xi)$, accompanied by $P \rightarrow P|h'_{,\xi}|$ preserves the Robinson-Trautman metric. Thus, locally, without loss of generality we can assume $h'(\xi) = \xi$. In this case substituting (B.12) into (B.4) yields

$$P^2 = \frac{c_3}{K_{,x}} \quad (\text{B.13})$$

where now $K = K(u, x)$ is considered as a function of u and $x = \text{Re } \xi$. Given (B.13) we see that $P = P(u, x)$ and equation (28) takes the form

$$c_3 P_{,x} - 12m P_{,u} = 0 \quad (\text{B.14})$$

It follows from (B.14) that P is a function of the variable $s = x + d(u)$, where

$$d_{,u} = \frac{c_3}{12m} . \quad (\text{B.15})$$

Equation (B.13) shows that

$$K = c_3(K'(s) + c_5(u)) \quad (\text{B.16})$$

$$P^2 = \frac{1}{K'_{,s}} , \quad (\text{B.17})$$

where K' and c_5 are functions of s and u , respectively. Substituting (B.16) and (B.17) into relation (5) implies that c_3 and c_5 are constants. Due to (B.15) and the remaining freedom of linear transformations of ξ we can assume that

$$K = K(x + u) , \quad P^2 = \frac{12m}{K_{,x}} . \quad (\text{B.18})$$

Now equation (B.6) takes the form

$$6mK_{,x} = -\frac{1}{3}K^3 + bK + c \quad (\text{B.19})$$

where b and c are constants. Robinson-Trautman metrics satisfying conditions (B.18) and (B.19) are equivalent to the C-metrics [11]. Thus, the Schwarzschild metric and the C-metric are the only vacuum Robinson-Trautman metrics which satisfy condition (26).

References

- [1] Robinson I., Trautman A., *Some spherical gravitational waves in general relativity*, Proc. Roy. Soc. A **265** 463 (1962)
- [2] Stephani H, Kramer D, MacCallum M A H, Hoenselaers C and Herlt E *Exact Solutions to Einstein's Field Equations, Second Edition*, (Cambridge: Cambridge University Press) (2003)
- [3] Chruściel P. T., *On the global structure of Robinson–Trautman spacetimes*, Proc. R. Soc. Lond. A **436** 299 (1992)
- [4] Ashtekar A., Krishnan B., *Isolated and Dynamical Horizons and Their Applications*, Living Rev. Relativity **7**, (2004), 10 [Online Article]: cited [July 18, 2008], <http://www.livingreviews.org/lrr-2004-10>
- [5] Tod P., *Analogues of the past horizon in Robinson–Trautman spacetimes*, Class. Quantum Grav. **8** 1159 (1989)
- [6] Chow E. W. M., Lun A. W.-C., *Apparent Horizons in Vacuum Robinson–Trautman Spacetimes*, J. Austral. Math. Soc. Ser. B **41** 217 (1999), <http://arxiv.org/abs/gr-qc/9503065> (1995)
- [7] Griffiths J. B., Krtouš P., and Podolský J., *Interpreting the C-metric*, Class. Quantum Grav. **23** 6745 (2006)
- [8] Andersson L., Mars M., Simon W., *Local Existence of Dynamical and Trapping Horizons*, Phys. Rev. Lett. **95**, 111102 (2005)
- [9] Korzyński M., *Isolated and dynamical horizons from a common perspective*, Phys. Rev. **D74**, 104029 (2006)
- [10] Kinnersley W., *Type D vacuum metrics*, J. Math. Phys. **10**, 1195 (1969)
- [11] Kinnersley W., *Field of an Arbitrarily Accelerating Point Mass*, Phys. Rev. **186**, 1335 (1969)